

②

AD-A212 618

REPORT DOCUMENTATION PAGE				
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY SEP 13 1989		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE B		4. PERFORMING ORGANIZATION REPORT NUMBER(S)		
5a. NAME OF PERFORMING ORGANIZATION Old Dominion University		5b. OFFICE SYMBOL (If applicable)		5. MONITORING ORGANIZATION REPORT NUMBER(S) ARO 24834.2-MA
6a. ADDRESS (City, State, and ZIP Code) Norfolk, VA 23508-0369		7a. NAME OF MONITORING ORGANIZATION U. S. Army Research Office		
6b. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211		7b. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U. S. Army Research Office		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAAL03-88-K-0076
8c. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211		10. SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. PROJECT NO. TASK NO. WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) Strong Moderate Deviation Theorems for m-Dependent Random Variables				
12. PERSONAL AUTHOR(S) Narasinga R. Changanthy				
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) August 1989
15. PAGE COUNT 9				
16. SUPPLEMENTARY NOTATION The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.				
17. COSATI CODES FIELD GROUP SUB-GROUP			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Deviation Theorems, Random Variables	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This document considers a stationary sequence $\{X_1, X_2, \dots\}$ of m-dependent random variables. Let $S_n = \sum_{i=1}^n X_i$ be the partial sum. Under some moment conditions, in this paper we obtain asymptotic expression for the probability of moderate deviations, $P(S_n > x_n)$, where $x_n = O(\sqrt{\log(n)})$. This result extends some well known results obtained for independent and identically distributed sequences of random variables. (KR)				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)	
			22c. OFFICE SYMBOL	

DEPARTMENT OF MATHEMATICAL SCIENCES
COLLEGE OF SCIENCES
OLD DOMINION UNIVERSITY
NORFOLK, VIRGINIA 23529

**STRONG MODERATE DEVIATION THEOREMS FOR m -DEPENDENT
RANDOM VARIABLES**

By

Narasinga R. Chaganty, Principal Investigator

Interim Technical Report
For the period ended June 30, 1989

Prepared for the
U.S. Army Research Office
Post Office Box 12211
Research Triangle Park, NC 27709-2211

Under
Contract DAAG03-88-K0076
Dr. Jagdish Chandra, Director
Mathematical Sciences Division

Submitted by the
Old Dominion University Research Foundation
Post Office Box 6369
Norfolk, Virginia 23508-0369

! August 1989

Strong Moderate Deviation Theorems for m-Dependent Random Variables†

By

Narasinga Rao Chaganty
Old Dominion University

Abstract

Consider a stationary sequence $\{X_1, X_2, \dots\}$ of m-dependent random variables. Let $S_n = \sum_{i=1}^n X_i$ be the partial sum. Under some moment conditions, in this paper we obtain asymptotic expression for the probability of moderate deviations, $P(S_n > x_n)$, where $x_n = O(\sqrt{\log(n)})$. This result extends some well known results obtained for independent and identically distributed sequences of random variables.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

†Research partially supported by the U.S. Army Research Office Grant number DAAL03-88-K-0076. The United States Government is authorized to reproduce and distribute reprints for Governmental purposes notwithstanding any copyright notation thereon.

AMS (1980) Subject classifications: Primary 60F10, Secondary 60G10.

Key words: Moderate Deviations, m-Dependence.

1. Introduction. Let $\{X_n, n \geq 1\}$ be a sequence of i.i.d. random variables with mean 0 and variance 1. Let $S_n = \sum_{i=1}^n X_i$ be the n^{th} partial sum. The theory of moderate deviations introduced by Rubin and Sethuraman (1965a) is concerned with obtaining asymptotic expression for

$$(1.1) \quad P(S_n > x_n)$$

where $x_n = O(\sqrt{\log(n)})$, under some moment conditions which are less restrictive than the assumption of finiteness of the moment generating function of X_1 . In a subsequent paper Rubin and Sethuraman (1965b) showed that the asymptotic expression for (1.1) is useful to compare test statistics via Bayes risk efficiency. We shall call a result which gives the asymptotic expression for (1.1) a weak moderate deviation result. On the other hand a strong moderate deviation theorem gives an asymptotic expression to the probability of the event $\{S_n > x\}$ which is valid uniformly in the interval $-A \leq x \leq c\sqrt{\log(n)}$. In this paper we obtain strong moderate deviation theorem for the partial sums of a stationary sequence $\{X_n, n \geq 1\}$ of m -dependent random variables. Note that a sequence $\{X_n, n \geq 1\}$ is said to be m -dependent if (X_1, \dots, X_r) and (X_s, X_{s+1}, \dots) are independent whenever $s - r > m$. The sequence is said to be stationary if $(X_{i+1}, \dots, X_{i+k})$ has the same distribution as $(X_{j+1}, \dots, X_{j+k})$ for all $k \geq 1$ and $i \neq j$.

2. Main Results. In this section we establish the main theorem of this paper. Theorem 2.1 below obtains a strong moderate deviation theorem for partial sums of m -dependent sequence of random variables.

Theorem 2.1. Consider a stationary sequence $\{X_n, n \geq 1\}$ of m -dependent random variables. Let $E(X_1) = 0$ and $\sigma^2 = \text{Var}(X_1) + 2 \sum_{j=1}^m \text{Cov}(X_1, X_{1+j})$ be finite. Let $S_n = \sum_{i=1}^n X_i$. If $E|X_1|^p < \infty$ for some $p > c^2 + 2$, where $c > 0$, then

$$(2.1) \quad P\left(\frac{S_n}{\sqrt{n}\sigma} > x\right) = [1 - \Phi(x)] \left[1 + o\left(\frac{1}{\log(n)}\right)\right]$$

uniformly in the region $-A \leq x \leq c\sqrt{\log(n)}$, where $A \geq 0$ is a constant and Φ denotes the distribution function of standard normal.

Vandemaele and Veraverbeke (1982) obtained strong moderate deviation theorems for L-statistics which are functions of independent and identically distributed sequences of random variables. A special case of their Theorem 1 yields the following Lemma 2.2. We will need Lemma 2.2 in the proof of Theorem 2.1.

Lemma 2.2. Let $\{X_n, n \geq 1\}$ be sequence of i.i.d. random variables with mean zero and variance σ^2 . Let $S_n = \sum_{i=1}^n X_i$ be the n^{th} partial sum. If $E|X_1|^p < \infty$ for some $p > c^2 + 2$, ($c > 0$) then (2.1) holds uniformly in the region $-A \leq x \leq c\sqrt{\log(n)}$, where $A \geq 0$ is a constant.

In the case $m = 0$, Theorem 2.1 yields Lemma 2.2 and in this sense our main result generalizes the result of Vandemaele and Veraverbeke (1982) to m -dependent random variables.

Proof of Theorem 2.1. We shall use the blocking technique used by Hoeffding and Robbins (1948) in proving central limit theorem for m -dependent random variables. Let $0 < \alpha < \min\{1, 1/p\}$ be fixed and $k = \max\{2m, [n^\alpha]\}$. We can write $n = k\nu + r$, where

$0 \leq r \leq k$. For n sufficiently large we can partition the n^{th} partial sum as follows:

$$\begin{aligned}
 S_n &= [X_1 + \dots + X_{k-m}] + [X_{k-m+1} + \dots + X_k] + [X_{k+1} + \dots + X_{2k-m}] \\
 &\quad + [X_{2k-m+1} + \dots + X_{2k}] + \dots + [X_{\nu k-m+1} + \dots + X_{\nu k}] \\
 &\quad + [X_{\nu k+1} + \dots + X_n]. \\
 (2.2) \quad &= U_{n1} + R_{n1} + U_{n2} + R_{n2} + \dots + U_{n\nu} + R_{n\nu} + T_{nr} \\
 &= [U_{n1} + \dots + U_{n\nu}] + [R_{n1} + R_{n2} + \dots + R_{n\nu}] + T_{nr} \\
 &= U_n + R_n + T_{nr}. \quad (\text{say}).
 \end{aligned}$$

Note that U_n is the sum of ν i.i.d. random variables and R_n is also the sum of ν i.i.d. random variables. Let $\delta_n = 1/(\log(n))^2$. It is easy to verify that the following important identity holds:

$$\begin{aligned}
 (2.3) \quad &P\left(\frac{U_n}{\sqrt{n}\sigma} > x + 2\delta_n\right) - P\left(\frac{R_n}{\sqrt{n}\sigma} < -\delta_n\right) - P\left(\frac{T_{nr}}{\sqrt{n}\sigma} < -\delta_n\right) \\
 &\leq P\left(\frac{S_n}{\sqrt{n}\sigma} > x\right) \\
 &\leq P\left(\frac{U_n}{\sqrt{n}\sigma} > x - 2\delta_n\right) + P\left(\frac{R_n}{\sqrt{n}\sigma} > \delta_n\right) + P\left(\frac{T_{nr}}{\sqrt{n}\sigma} > \delta_n\right).
 \end{aligned}$$

Since $[1 - \Phi(x)]^{-1} = O((c\sqrt{\log(n)})n^{c^2/2})$ uniformly in $-A \leq x \leq c\sqrt{\log(n)}$, ($A > 0, c > 0$), the proof of the theorem will be complete once we establish the following Lemma 2.3.

Lemma 2.3. *Let U_n, R_n and T_{nr} be as defined above. Then under the hypothesis of Theorem 2.1 we have the following:*

$$\begin{aligned}
 (A) \quad &P\left(\frac{U_n}{\sqrt{n}\sigma} > x \pm 2\delta_n\right) = [1 - \Phi(x)] \left[1 + o\left(\frac{1}{\log(n)}\right)\right] \\
 &\text{uniformly in } -A \leq x \leq c\sqrt{\log(n)}.
 \end{aligned}$$

$$(B) \ P \left(\left| \frac{R_n}{\sqrt{n}\sigma} \right| > \delta_n \right) = o \left((\log(n))^{-3/2} n^{-c^2/2} \right).$$

$$(C) \ P \left(\left| \frac{T_{nr}}{\sqrt{n}\sigma} \right| > \delta_n \right) = o \left((\log(n))^{-3/2} n^{-c^2/2} \right).$$

Proof of (A). Note that $U'_n = \frac{U_n}{k}$ is the sum of ν i.i.d. random variables with mean zero and variance given by

$$(2.4) \quad M^2 = \frac{1}{k^2} [(k-m) \text{Var}(X_1) + 2 \sum_{j=1}^m (k-m-j) \text{Cov}(X_1, X_{1+j})].$$

Now,

$$(2.5) \quad \begin{aligned} P \left(\frac{U_n}{\sqrt{n}\sigma} > x \right) &= P \left(\frac{\sqrt{\nu}kM}{\sqrt{n}\sigma} \frac{U'_n}{M\sqrt{\nu}} > x \right) \\ &= P \left(\frac{U'_n}{M\sqrt{\nu}} > xc_n \right) \end{aligned}$$

where

$$(2.6) \quad \begin{aligned} c_n &= \frac{\sqrt{n}\sigma}{M\sqrt{\nu}k} = \left[\frac{\sigma^2}{kM^2} \right]^{1/2} \left[\frac{n}{k\nu} \right]^{1/2} \\ &= \left[\frac{\sigma^2}{kM^2} \right]^{1/2} \left[1 + \frac{r}{k\nu} \right]^{1/2}. \end{aligned}$$

Note that

$$(2.7) \quad \begin{aligned} kM^2 &= \frac{1}{k} [(k-m) \text{Var}(X_1) + 2 \sum_{j=1}^m (k-m-j) \text{Cov}(X_1, X_{1+j})] \\ &= \sigma^2 - \frac{1}{k} [m \text{Var}(X_1) + 2 \sum_{j=1}^m (m+j) \text{Cov}(X_1, X_{1+j})] \\ &= \sigma^2 + O(n^{-\alpha}) = \sigma^2 [1 + O(n^{-\alpha})] \end{aligned}$$

since $k = O(n^\alpha)$. Thus we get

$$(2.8) \quad \begin{aligned} c_n &= [1 + O(n^{-\alpha})]^{1/2} \left(1 + \frac{r}{k\nu} \right)^{1/2} \\ &= [1 + O(n^{-\beta})] \quad \text{for some } \beta > 0. \end{aligned}$$

Therefore we have shown that

$$P\left(\frac{U_n}{\sqrt{n}\sigma} > x\right) = P\left(\frac{U'_n}{M\sqrt{\nu}} > c_n x\right)$$

where $c_n = [1 + O(n^{-\beta})]$. Hence,

$$\begin{aligned} P\left(\frac{U_n}{\sqrt{n}\sigma} > x \pm \frac{2}{(\log(n))^2}\right) &= P\left(\frac{U'_n}{M\sqrt{\nu}} > c_n \left[x \pm \frac{2}{(\log(n))^2}\right]\right) \\ (2.10) \qquad \qquad \qquad &= P\left(\frac{U'_n}{M\sqrt{\nu}} > x_n\right) \end{aligned}$$

where

$$\begin{aligned} x_n &= c_n \left[x \pm \frac{2}{(\log(n))^2}\right] \\ (2.11) \qquad \qquad \qquad &= [1 + O(n^{-\beta})] \left[x \pm \frac{2}{(\log(n))^2}\right]. \end{aligned}$$

Let δ be such that $\sqrt{p-2} > c + \delta$. Then applying Lemma 2.2 with c replaced by $c + \delta$ we get

$$(2.12) \qquad P\left(\frac{U'_n}{M\sqrt{\nu}} > x_n\right) = [1 - \Phi(x_n)] \left[1 + o\left(\frac{1}{\log(n)}\right)\right]$$

uniformly in the interval $-A \leq x_n \leq (c + \delta)\sqrt{\log(n)}$. Also using Lemma A1 of Vandermale and Veraverbeke (1982) and (2.11) we get

$$(2.13) \qquad [1 - \Phi(x_n)] = [1 - \Phi(x)] \left[1 + o\left(\frac{1}{\log(n)}\right)\right].$$

Combining (2.10), (2.12) and (2.13) we have

$$P\left(\frac{U_n}{\sqrt{n}\sigma} > x \pm \frac{2}{(\log(n))^2}\right) = [1 - \Phi(x)] \left[1 + o\left(\frac{1}{\log(n)}\right)\right]$$

uniformly in $-A \leq x \leq c\sqrt{\log(n)}$. This completes the proof of (A).

Proof of (B). Note that R_n is the sum of ν i.i.d. random variables with mean zero and variance

$$(2.14) \quad \sigma_1^2 = m \text{Var}(X_1) + 2 \sum_{j=1}^{m-1} (m-j) \text{Cov}(X_1, X_{1+j}).$$

Fix constants α and c_1 such that $\alpha < 1 - \frac{c^2}{p-2}$, $c_1^2 > \frac{c^2}{(1-\alpha)}$ and $p > 2 + c_1^2$. Applying Theorem 1 of Rubin and Sethuraman (1965a) we get for sufficiently large n ,

$$(2.15) \quad \begin{aligned} P \left(\left| \frac{R_n}{\sqrt{n}\sigma} \right| > \delta_n \right) &= P \left(\left| \frac{R_n}{\sqrt{\nu}} \right| > \frac{\sigma\sqrt{n}}{\sqrt{\nu}(\log(n))^2} \right) \\ &\leq P \left(\left| \frac{R_n}{\sqrt{\nu}} \right| > \sigma_1 c_1 \sqrt{\frac{\log(\nu)}{\nu}} \right) \\ &\sim \frac{\nu^{-c_1^2/2}}{c_1 \sqrt{2\pi \log(\nu)}} \end{aligned}$$

since $p > c_1^2 + 2$. Using the fact $\nu = O(n^{1-\alpha})$ and $(1-\alpha)c_1^2 > c^2$ and (2.15), we get that

$$P \left(\left| \frac{R_n}{\sqrt{n}\sigma} \right| > \delta_n \right) = o \left((\log(n))^{-3/2} n^{-c^2/2} \right).$$

This proves (B).

Proof of (C). Applying Chebyshev's inequality we get

$$(2.16) \quad \begin{aligned} P \left(\left| \frac{T_{nr}}{\sqrt{n}\sigma} \right| > \delta_n \right) &\leq \frac{E|T_{nr}|^p}{(\sigma\sqrt{n}\delta_n)^p} \\ &\leq \text{const.} \frac{k^p}{(\sigma\sqrt{n}\delta_n)^p}. \end{aligned}$$

Note that $k = O(n^\alpha)$ and hence

$$\begin{aligned}
(2.17) \quad n^{c^{2/2}} (\log(n))^{3/2} \frac{k^p}{(\sqrt{n}\delta_n)^p} &= \frac{n^{c^{2/2}} n^{p\alpha}}{n^{p/2}} (\log(n))^{2p+3/2} \\
&= n^{-\frac{1}{2}(p-2p\alpha-c^2)} (\log(n))^{2p+3/2} \\
&\rightarrow 0 \quad \text{as } n \rightarrow \infty,
\end{aligned}$$

since $0 < \alpha < \frac{1}{p}$ and $p > c^2 + 2$. The proof of (C) now follows combining (2.16) and (2.17).

Remark 2.4. Theorem 2.1 suggests that it is possible to obtain strong moderate deviation theorems for L-statistics and U-statistics which are functions of m-dependent sequences of random variables in the same spirit as Vandemaële and Veraverbeke(1982) and Ghosh(1974).

3. References.

1. Ghosh, M.(1974). Probabilities of moderate deviations under m-dependence. *The Canadian Journal of Statistics*, **2**, 157-168.
2. Rubin, H. and Sethuraman, J.(1965a). Probabilities of moderate deviations. *Sankhyā*, **A**, **27**, 325-346.
3. Rubin, H. and Sethuraman, J.(1965b). Bayes risk efficiency. *Sankhyā*, **A**, **27**, 347-356.
4. Vandermaele, M. and Veraverbeke, N.(1982). Cramér type large deviations and moderate deviations for linear combinations of order statistics. *Ann. Probab.*, **10**, 423-434.